

ItMM 5 Functions

5.1 Which of the following relations are functions? Find their domains and sets of values. Which of them are injections?

- a) for $x, y \in \mathbb{R}$, $xRy \Leftrightarrow x^3 = y^4$,
- b) for $x, y \in \mathbb{R}$, $xRy \Leftrightarrow \frac{y-1}{x} = 1$,
- c) for $x, y, z \in \mathbb{R}$, $(x, y)Sz \Leftrightarrow x + y + z^2 = 1$
- d) for $x, y \in \mathbb{N}$, $xUy \Leftrightarrow x$ is the greatest prime divisor of y ,
- f) for polynomial p and $x \in \mathbb{R}$, $pTx \Leftrightarrow p(x) = 0$,
- g) for $A, B \subseteq X$, $AFB \Leftrightarrow A \cup B = X$ and $A \cap B = \emptyset$,
- h) for $A \subseteq \mathbb{N}$, $x \in \mathbb{N}$, $AGx \Leftrightarrow x$ is a product of all elements of A ,
- i) for a quadratic polynomial f (with real coefficients) and $A \subseteq \mathbb{R}$, $fVA \Leftrightarrow (\forall x \in \mathbb{R})x \in A \Leftrightarrow f(x) = 1$,
- j) for $x, y \in \mathbb{R}$, $xVy \Leftrightarrow (\exists z \in \mathbb{R})y = \frac{x+z}{2}$

5.2 For a function f and a subset A of its domain find $f(A)$ and $f^{-1}(f(A))$

- a) $f(x) = x^2$ for $x \in \mathbb{R}$, $A = [-2; 3)$,
- b) $f(x) = \cos x$ for $x \in \mathbb{R}$, $A = [-\pi/2; \pi/2]$,
- c) $f(x, y) = x \cdot y$ for $x, y \in \mathbb{N}$, $A = [0, \infty) \times [0, \infty)$
- d) $f(x, y) = x^2 - y^2$ for $x, y \in \mathbb{N}$, $A = [0, \infty) \times [0, \infty)$.
- e) $f(x, y) = x + y$ for $x, y \in \mathbb{R}$, $A = \{(0, 0), (0, 1)\}$.
- f) $f(x, y) = \frac{x}{y+1}$ for $x, y \in \mathbb{N}$, $A = \{(1, 2)\}$.
- g) $f(x, y) = (x + y, x - y)$ for $x, y \in \mathbb{R}$, $A = \{(x, y) \in \mathbb{R}^2 : x = y\}$.
- h) $f(x, y) = x^2 + y^2$ for $x, y \in \mathbb{R}$, $A = \{(x, y) \in \mathbb{R}^2 : 1 < x < 2 \wedge 0 < y < 1\}$.
- i) $f(x, y) = \max(x, y)$ for $x, y \in \mathbb{R}$, i) $A = [0, 1] \times [0, 1]$, ii) $A = \{(x, y) \in \mathbb{R}^2 : x^2 + y^2 = 1\}$.
- j) $f(p) = p(1)$ for a polynomial p with real coefficients, A is the family of all linear functions of the form $ax - a$.
- k) $f(n) =$ sum of all prime divisors of n for $n \in \mathbb{N}$, $A = \{4, 6\}$,
- l)* $f(X) = \{x \in \mathbb{R} : (\exists y)(x, y) \in X\}$ for $X \subseteq \mathbb{R}^2$, A is the family of all singletons.

5.3 Let $A \subseteq X$ and $\chi_A : X \rightarrow \{0, 1\}$ be a characteristic function of a set A i.e. for $x \in X$, $\chi_A(x) = 1$ for $x \in A$ and $\chi_A(x) = 0$ for $x \notin A$. Prove that the function F defined $F(A) = \chi_A$ is a on-to-on mapping of the powerset of X onto set $\{0, 1\}^X$.

5.4 Let $A, B \subseteq X$ and $x \in X$. Prove

- a) $\chi_{-A}(x) = 1 - \chi_A(x)$,
- b) $\chi_{A \cap B}(x) = \chi_A(x) \cdot \chi_B(x)$.